

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Association of menopausal characteristics and risk of coronary heart disease: a pan-European case-cohort analysis**

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1704686> since 2020-02-28T16:46:59Z

*Published version:*

DOI:10.1093/ije/dyz016

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**Genetically determined reproductive aging and cardiovascular risk factors and coronary heart disease risk: a two-sample Mendelian Randomization study**

Veerle Dam<sup>1,2</sup>, N Charlotte Onland-Moret<sup>\*1</sup>, Stephen Burgess<sup>3,4,5</sup>, Maria-Dolores Chirlaque<sup>6,33</sup>, Sanne A E Peters<sup>1,7</sup>, Ewoud Schuit<sup>1</sup>, Kaja Tikk<sup>8,9</sup>, Elisabete Weiderpass<sup>10,11,12,13</sup>, Clare Oliver-Williams<sup>4,5</sup>, Angela M Wood<sup>4</sup>, Anne Tjønneland<sup>14,15</sup>, Christina C Dahm<sup>16</sup>, Kim Overvad<sup>16,17</sup>, Francoise Clavel-Chapelon<sup>18</sup>, Heiner Boeing<sup>19</sup>, Antonia Trichopoulou<sup>20,21</sup>, Pietro Ferrari<sup>22</sup>, Giovanna Masala<sup>23</sup>, Claudia Agnoli<sup>24</sup>, Rosario Tumino<sup>25</sup>, Giuseppe Matullo<sup>26,27</sup>, Salvatore Panico<sup>28</sup>, Jolanda M A Boer<sup>29</sup>, W M Monique Verschuren<sup>1,29</sup>, Marit Waaseth<sup>30</sup>, Virginia Menéndez<sup>31</sup>, Miguel Rodríguez-Barranco<sup>32,33</sup>, Olatz Mokoroa<sup>33,34</sup>, Conchi Moreno-Iribas<sup>35</sup>, Olle Melander<sup>36</sup>, Sophia Harlid<sup>37</sup>, Maria Nordendahl<sup>38</sup>, Timothy J Key<sup>39</sup>, Elio Riboli<sup>40</sup>, Carmen Santiuste<sup>33,41</sup>, Tilman Kühn<sup>42</sup>, Verena Katzke<sup>42</sup>, Claudia Langenberg<sup>43</sup>, Nicholas J Wareham<sup>43</sup>, Heribert Schunkert<sup>44,45</sup>, Jeanette Erdmann<sup>46</sup>, Christina Willenborg<sup>46</sup>, Christian Hengstenberg<sup>47</sup>, Marcus E Kleber<sup>48</sup>, Graciela Delgado<sup>48</sup>, Winfried März<sup>48,49,50</sup>, Stavroula Kanoni<sup>51</sup>, George Dedoussis<sup>52</sup>, Panos Deloukas<sup>51,53,54</sup>, Majid Nikpay<sup>55</sup>, Ruth McPherson<sup>55</sup>, Markus Scholz<sup>56,57</sup>, Andrej Teren<sup>57,58</sup>, Adam S Butterworth<sup>4</sup>, Yvonne T van der Schouw<sup>1</sup>

<sup>\*</sup>Corresponding author

<sup>1</sup>Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands

<sup>2</sup>Netherlands Heart Institute, Utrecht, the Netherlands

<sup>3</sup>MRC Biostatistics Unit, University of Cambridge, Cambridge, United Kingdom

21 <sup>4</sup> MRC/BHF Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care,  
22 University of Cambridge, Cambridge, United Kingdom

23 <sup>5</sup>Homerton College, Hills Rd, Cambridge, United Kingdom

24 <sup>6</sup>Department of Epidemiology, Regional Health Authority, IMIB-Arrixaca, Murcia University,  
25 Murcia, Spain

26 <sup>7</sup>The George Institute for Global Health, University of Oxford, Oxford, United Kingdom

27 <sup>8</sup>Division of Clinical Epidemiology and Aging Research, German Cancer Research Centre  
28 (DKFZ), Heidelberg, Germany

29 <sup>9</sup>German Cancer Consortium (DKTK), German Cancer Research Centre (DKFZ), Heidelberg,  
30 Germany

31 <sup>10</sup>Department of Community Medicine, Faculty of Health Sciences, University of Tromsø, The  
32 Arctic University of Norway, Tromsø, Norway

33 <sup>11</sup>Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm,  
34 Sweden

35 <sup>12</sup>Genetic Epidemiology Group, Folkhälsan Research Center and Faculty of Medicine, University  
36 of Helsinki, Helsinki, Finland

37 <sup>13</sup>Department of Research, Cancer Registry of Norway, Institute of Population-Based Cancer  
38 Research, Oslo, Norway

39 <sup>14</sup>Danish Cancer Society Research Center, Copenhagen, Denmark

40 <sup>15</sup>Department of Public Health, Faculty of Health and Medical Sciences, University of  
41 Copenhagen, Copenhagen, Denmark

42 <sup>16</sup>Department of Public Health, Aarhus University, Denmark

43 <sup>17</sup>Department of Cardiology, Aalborg University Hospital, Denmark

44 <sup>18</sup>INSERM, Centre for Research in Epidemiology and Population Health (CESP), U1018,  
45 Nutrition, Hormones, and Women's Health Team, Institut Gustave Roussy, Villejuif, France

46 <sup>19</sup>Department of Epidemiology, German Institute of Human Nutrition (DIfE), Potsdam-  
47 Rehbrücke, Germany

48 <sup>20</sup>Hellenic Health Foundation, Athens, Greece

49 <sup>21</sup>WHO Collaborating Center for Nutrition and Health, Unit of Nutritional Epidemiology and  
50 Nutrition in Public Health, Dept. of Hygiene, Epidemiology and Medical Statistics, School of  
51 Medicine, National and Kapodistrian University of Athens, Greece.

52 <sup>22</sup>International Agency for Research on Cancer, Lyon, France

53 <sup>23</sup>Cancer Risk Factors and Life-Style Epidemiology Unit, Institute for Cancer Research,  
54 Prevention and Clinical Network - ISPRO, Florence, Italy

55 <sup>24</sup>Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan,  
56 Italy

57 <sup>25</sup>Cancer Registry and Histopathology Department, 'Civic - M.P. Arezzo' hospital, ASP Ragusa,  
58 Italy

- 59 <sup>26</sup>Department of Medical Sciences, University of Torino
- 60 <sup>27</sup>Italian Institute for Genomic Medicine –IIGM/HuGeF, Italy
- 61 <sup>28</sup>Dipartimento di medicina clinica e chirurgia, Federico II University, Naples, Italy
- 62 <sup>29</sup>National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands
- 63 <sup>30</sup>Department of Pharmacy, Faculty of Health Sciences, UiT the Arctic University of Norway,
- 64 Tromsø, Norway
- 65 <sup>31</sup>Public Health Directorate, Asturias, Spain
- 66 <sup>32</sup>Escuela Andaluza de Salud Pública. Instituto de Investigación Biosanitaria ibs.GRANADA,
- 67 Universidad de Granada, Granada, Spain
- 68 <sup>33</sup>CIBER de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain
- 69 <sup>34</sup>Public Health Division of Gipuzkoa, Biodonostia Research Institute, San Sebastian, Spain
- 70 <sup>35</sup>Instituto de Salud Pública de Navarra, IdiSNA, Navarre Institute for Health Research,
- 71 REDISSEC Pamplona, Spain
- 72 <sup>36</sup>Department of Clinical Sciences, Malmö, Lund University, Malmö
- 73 <sup>37</sup>Department of Radiation Sciences, Oncology, Umea University, Umea, Sweden
- 74 <sup>38</sup>Department of Public Health and Clinical Medicine, Umea University, Umea, Sweden
- 75 <sup>39</sup>Nuffield Department of Population Health, University of Oxford, Oxford, England

76 <sup>40</sup>Department of Epidemiology and Biostatistics, School of Public Health, Imperial College  
77 London, London, United Kingdom

78 <sup>41</sup>Department of Epidemiology, Murcia Regional Health Authority, IMIB-Arrixaca, Murcia,  
79 Spain

80 <sup>42</sup>Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Foundation under  
81 Public Law, Heidelberg, Germany

82 <sup>43</sup>MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge,  
83 United Kingdom

84 <sup>44</sup>Deutsches Herzzentrum München, Technische Universität München, Munich, Germany

85 <sup>45</sup>DZHK (German Center for Cardiovascular Research), partner site Munich Heart Alliance,  
86 Munich, Germany

87 <sup>46</sup>Institute for Cardiogenetics, University of Lübeck, Lübeck, Germany

88 <sup>47</sup>Department of Internal Medicine II, Division of Cardiology, Medical University of Vienna,  
89 Vienna, Austria

90 <sup>48</sup>Vth Department of Medicine, Medical Faculty Mannheim, Heidelberg University, Mannheim,  
91 Germany

92 <sup>49</sup>Synlab Academy, Synlab Holding Deutschland GmbH, Mannheim, Germany

93 <sup>50</sup>Clinical Institute of Medical and Chemical Laboratory Diagnostics, Medical University of  
94 Graz, Graz, Austria

95 <sup>51</sup>William Harvey Research Institute, Barts and The London School of Medicine and Dentistry,  
96 Queen Mary University of London, London EC1M 6BQ, United Kingdom

97 <sup>52</sup>Department of Nutrition-Dietetics/Harokopio University, Athens, Greece

98 <sup>53</sup>Centre for Genomic Health, Queen Mary University of London, London EC1M 6BQ, United  
99 Kingdom

100 <sup>54</sup>Princess Al-Jawhara Al-Brahim Centre of Excellence in Research of Hereditary Disorders  
101 (PACER-HD), King Abdulaziz University, Jeddah 21589, Saudi Arabia

102 <sup>55</sup>Ruddy Canadian Cardiovascular Genetics Centre, University of Ottawa Heart Institute, Ottawa,  
103 Canada

104 <sup>56</sup>Institute for Medical Informatics, Statistics and Epidemiology, University of Leipzig, Germany

105 <sup>57</sup>LIFE Research Center for Civilization Diseases, University of Leipzig, Leipzig, Germany

106 <sup>58</sup>Heart Center Leipzig, Leipzig, Germany

107

108 Corresponding author:

109 Dr. N. Charlotte Onland-Moret

110 [n.c.onland@umcutrecht.nl](mailto:n.c.onland@umcutrecht.nl)

111 Huispostnummer STR. 6.131

112 P.O. Box 85500

113 3508 GA Utrecht

114 the Netherlands

115

116

117



## Abstract

**Background:** Accelerated reproductive aging, in women indicated by early natural menopause, is associated with an increased risk of coronary heart disease (CHD) in observational studies. Genomic variants for age at natural menopause (ANM) have been implicated in genome stability, immune function and mitochondrial biogenesis, which are not sex-specific processes. We aimed to establish the causal association between reproductive aging and (non-)fatal CHD and CHD risk factors using ANM variants as a measure for genetically determined reproductive aging in women and in men, since genome-wide association studies (GWAS) for reproductive aging traits in men are lacking.

**Methods:** We performed a 2-sample Mendelian Randomization (MR) using four methods: the simple median-based method, the weighted median-based method, the standard inverse-variance weighted (IVW) regression and the MR-Egger regression. Summary statistics were pooled from three studies with together 417,579 participants from European descent, including 49,150 CHD cases. Publicly available GWAS and EPIC-CVD were pooled for total cholesterol, high density lipoprotein cholesterol, triglycerides, HbA1c, and glucose.

**Results:** Our MR analyses show no association between genetically determined reproductive aging and CHD risk in women (Relative Risk Estimate (RRE)<sub>IVW</sub>=0.99, 95% confidence interval (CI): 0.97;1.01), or any of the CHD risk factors. No associations were found in men.

**Conclusion:** Reproductive aging is not causally associated with CHD risk or CHD risk factors in women, nor in men. The association between early menopause and CHD risk in observational studies might be the result of residual confounding, reverse causation, or reflect a shared aetiology that results in both earlier menopause and higher CHD risk.

**Keywords:** Reproductive aging, Mendelian randomization, coronary heart disease,  
cardiovascular risk factors

**Key messages:**

- Genetically determined reproductive aging is not associated with coronary heart disease in women.
- Genetically determined reproductive aging is not associated with coronary heart disease in men, although the validity of the genetic instrument is not established in men.
- Genetically determined reproductive aging is not associated with cardiovascular risk factors (total cholesterol, high density lipoprotein cholesterol, triglycerides, apolipoprotein A1, apolipoprotein B, C-reactive protein, glucose and HbA1c).

## Introduction

Cardiovascular disease (CVD) is the leading cause of death in both men and women(1). Accelerated reproductive aging, as indicated by early menopause in women, has been associated with increased risk of CVD(2–5). The mechanisms underlying these associations are not fully understood yet; deterioration of traditional CVD risk factors, in particular cholesterol, has been suggested to play a role(6,7). Although men do not experience an abrupt start or stop of their reproductive period, there is limited evidence that in men reproductive functions, such as erectile dysfunction, sperm motility and morphology, and semen volume, also decline with aging(8–10). Some of these, e.g. erectile dysfunction and decreasing testosterone levels, sometimes referred to as andropause, have been associated with increased CVD risk as well(11,12). Since male reproductive aging is a gradual process into old age, it is more complicated to study health effects of accelerated reproductive aging in males.

In observational studies, it is difficult to disentangle the potential independent effect of accelerated reproductive aging on CVD risk from the effect of general aging, as residual confounding can still be present. Furthermore, reversed causality can also play a role here, as women with an unfavourable CVD risk profile have been reported to experience accelerated reproductive aging(13). Mendelian Randomization (MR) designs, exploiting the principle of random independent segregation of alleles at meiosis, are a means to establish causality in situations where randomized clinical trials are impossible(14,15). In MR studies, single nucleotide polymorphisms (SNPs) associated with the exposure as found in genome-wide association studies (GWAS) are used as instrumental variables.

To date, GWAS have been conducted for the reproductive aging trait age at natural menopause (ANM) in women, while GWAS for male reproductive aging traits are not available.

The ANM GWAS reported 56 SNPs that are mainly implicated in genome stability (DNA repair), immune function and mitochondrial biogenesis(16). As these mechanisms are not specific for women, we hypothesized that these mechanisms underlie reproductive aging in men as well.

A recent study in three cohorts suggested a harmful effect of ANM, genetically determined by the 56 SNPs, on CVD and CHD risk in women, but not in men. However, the sample size was small. Replication in a large sample size using publicly available data, conducted in women only, gave a null finding (17). This study did not investigate cardiovascular risk factors as an outcome.

The aims of the present study are to establish the causal association between reproductive aging and fatal or non-fatal CHD, and to gain more insight in possible mechanisms underlying the association between genetically determined reproductive aging and cardiovascular risk factors in women, using 56 SNPs associated with earlier ANM. Furthermore, we aim to establish whether the same mechanisms are associated with CHD and traditional cardiovascular risk factors in men as well. We used the same 56 ANM variants as a measure for genetically determined reproductive aging in men, postulating common genetic mechanisms of reproductive aging.

## Methods

### *Study populations and outcomes*

#### Fatal or non-fatal CHD

We used data from 417,579 participants of European descent (including 49,150 CHD cases) from three studies: the UK Biobank(18), a modified version of the CARDIoGRAMplusC4D consortium (m-CARDIoGRAMplusC4D) since we could only include those studies that provided us with sex-specific summary data (Cardiogenics, Thiseas, AMC-PAS, Duke 2, CCGB 2, ITH 2, OHGS A2, OHGS B2, OHGS C2, Germifs I, Germifs II, Germifs III, Germifs IV, LIFE-Heart and LURIC(19)), and the EPIC-CVD case-cohort study(20). Details of the three studies (UK Biobank, m-CARDIoGRAMplusC4D and EPIC-CVD), including definitions of fatal or non-fatal CHD in each study, can be found in supplement 1.

#### Traditional CHD risk factors

For the associations between genetically determined reproductive aging and CHD risk factors, we again used data from EPIC-CVD and combined these with publicly available GWAS summary statistics of the Global Lipids Genetics Consortium(21) (total cholesterol, high density lipoprotein (HDL) cholesterol, triglycerides) and MAGIC(22,23) (HbA1c, fasting glucose). Details on these consortia can be found in supplement 1. We did not have access to sex-specific data for these risk factors. Therefore, we could only perform a pooled MR analyses for men and women combined.

#### *Genotyping and SNP selection*

Genotyping in the UK Biobank was performed using the Affymetrix UK BiLEVE Axiom array and the Affymetrix UK Biobank Axiom Array(18,24). The m-CARDIoGRAMplusC4D studies have used various genotyping methods as described previously(19). EPIC-CVD participants were genotyped with the Human Core Exome array, Illumina 660 Quad array, and Omni Exome Express array. The Global Lipids Genetics Consortium and MAGIC also used different assays as described previously(21–23).

A recent genome-wide meta-analysis identified 56 SNPs associated with younger ANM among European descendants, 54 common HapMap SNPs and two Exome chip SNPs(16). All SNPs passed the threshold of  $p < 5e-6$ , but not all the threshold of  $p < 5e-8$ . No linkage disequilibrium (LD) at  $R^2 > 0.9$  was present among these 56 SNPs. Pleiotropic effects were investigated by searching the NHGRI-EBI GWAS Catalog(25) and Phenoscanner(26) for the SNPs or their proxies ( $R^2 > 0.8$ ). We used the 56 ANM variants as a measure for genetically determined reproductive aging in both women, and in men, since GWAS for reproductive aging traits in men are lacking.

### *Statistical analyses*

We verified whether the ANM variants were a valid instrument for the MR analysis in women by calculating the F-statistic according to the method described previously(27), using the SD (5.8 years) for ANM from the imputed data in the EPIC-CVD subcohort and the beta's for the ANM variants from the GWAS(16).

Regarding the outcome CHD, for UK Biobank and m-CARDIoGRAMplusC4D, odds ratios and standard errors for the SNP-CHD relations were derived through contact persons. For

EPIC-CVD, Prentice-weighted Cox proportional hazards regression adjusted for age, country, the first two principal components and array was used to calculate hazard ratios and standard errors for the EPIC-CVD case-cohort set. Regarding CHD risk factors, we derived effect estimates and standard errors for the cardiovascular risk factors (Global Lipids Genetics Consortium(21) for total cholesterol, HDL cholesterol and triglycerides, and MAGIC(22,23) for HbA1c and fasting glucose) using Phenoscanner(26). In the random subcohort of EPIC-CVD, we first imputed the missing observational data of EPIC-CVD (non-genetic data only) using multiple imputation with the MICE package in R(28) with 10 imputations and 50 iterations, including the CVD risk factors, SNPs and other baseline characteristics as predictors. Subsequently, we derived regression coefficients with linear regression in the subcohort only, separately in each imputation, using the same adjustments as for CHD. Thereafter we pooled the results with Rubin's Rule(29).

We performed a 2-sample MR using four separate methods to estimate causal effects for binary (CHD) and continuous (total cholesterol, HDL cholesterol, triglycerides, apolipoprotein A (apoA1), apolipoprotein B (apoB), C-reactive protein (CRP), glucose and HbA1c) outcomes: the simple median-based method, the weighted median-based method, the standard inverse-variance weighted (IVW) regression and the MR-Egger regression using the 'Mendelian Randomization' package in R(30). The IVW provides a consistent estimate and assumes that all assumptions of the instrumental variable are met, the median based and MR-Egger methods provide estimates under weaker assumptions, with the MR-Egger additionally providing an intercept that represents the average pleiotropic effect(31,32). When unbalanced horizontal pleiotropy is absent, results of all methods are expected to be consistent(33). We first conducted sex-specific MR analyses for CHD in all three studies (UK Biobank, m-CARDIoGRAMplusC4D, EPIC-

260 CVD) separately. Subsequently, we pooled the estimates with a fixed effect model as is standard  
261 in MR studies. Similarly, MR analyses were performed for each cardiovascular risk factor in  
262 each study separately (EPIC-CVD, Global Lipids Genetics Consortium, MAGIC) and then  
263 pooled using a fixed effects model. Sex-specific analyses were possible in EPIC-CVD only,  
264 therefore we pooled the results for both sexes for combining with Global Lipids Genetics  
265 Consortium and MAGIC). All analyses were conducted with R version 3.2.0(34).

266



## Results

Table 1 provides an overview of the numbers of cases and non-cases in UK Biobank, m-CARDIoGRAMplusC4D, and EPIC-CVD.

(Table 1 here)

The F-statistic for genetically determined reproductive aging in women was 93.7. Table 2 shows the results for the association between genetically determined reproductive aging and CHD per MR method stratified by sex and by study (UKBiobank, m-CARDIoGRAMplusC4D, and EPIC-CVD). In women, the IVW analyses in each study separately showed no causal association between genetically determined reproductive aging and CHD, nor when studies were pooled together (Relative Risk Estimate[RRE]<sub>IVW</sub>=0.99; 95% confidence interval [CI]=0.97;1.01). The MR-Egger method indicated no pleiotropic effects (intercept=0.004, p=0.318) and resulted in an RRE of 0.97 (95%CI=0.94;1.02) in the pooled data. Similar results were found for men with a pooled RRE<sub>IVW</sub> of 1.00 (95%CI=0.97;1.02), also indicating no pleiotropic effects (RRE<sub>MR-Egger</sub>=1.00 (95%CI=0.95;1.05), intercept=0.000, p=0.948).

(Table 2 here)

Table 3 shows the IVW results for the association between genetically determined reproductive aging and cardiovascular risk factors, with sex-specific estimates only from the EPIC-CVD subcohort and the sex-combined pooled estimates from both publicly available GWAS data and the EPIC-CVD subcohort. For each one-year decrease in genetically determined reproductive

aging, total cholesterol levels decreased with 0.025 mmol/L in women in IVW-analysis, however this was not statistically significant (95%CI= -0.056;0.005). Similarly, genetically determined reproductive aging was not causally associated with total cholesterol in men ( $\beta_{IVW}=0.024$  mmol/L, 95%CI= -0.011;0.059), nor in the pooled sex-combined results (pooled  $\beta_{IVW}=-0.005$  mmol/L, 95%CI= -0.007;0.017). Again, no pleiotropic effects were detected (supplement 2). Furthermore, no causal association was found for HDL cholesterol, triglycerides, ApoA1, ApoB, CRP, glucose, and HbA1c (table 3).

(Table 3 here)

## Discussion

This study did not find a causal association between reproductive aging and CHD risk or CHD risk factors, including cholesterol levels, in women. Furthermore, this study does not provide evidence for a causal association between reproductive aging and CHD risk or CHD risk factors in men.

Strengths of this study are that, to the best of our knowledge, this is the largest MR study of associations between reproductive aging and CHD to date with 20,169 CHD events in women and 27,397 in men. We used several methods for MR analyses all yielding consistent results for the tested hypotheses, and in women the instrument we used was strong (F-statistic 93.7). Some limitations need to be acknowledged. First, we cannot establish whether the ANM risk score is a valid instrument for reproductive aging in men. The F-statistic is calculated using observed menopausal age in women, but men do not have a similar trait with an abrupt stop in reproductive potential. Since the SNPs we used are mainly implicated in mechanisms that are not specific for women, we hypothesized that there are common mechanisms of reproductive aging for women and men, and that, therefore, the same variants can be used as marker for genetically determined reproductive aging in men. However, it needs to be acknowledged that corresponding phenotypic traits in men need to be further investigated. Second, the GWAS on ANM included women with an ANM between 40 and 60 years only and therefore did not include women with an extremely early menopause (<40) or premature ovary insufficiency (POI). Most of the observational studies did include women with an extremely early menopause or POI, and two recent systematic reviews and meta-analyses of observational studies showed that POI is associated with both fatal and non-fatal CHD and CVD(35,36). Although we could not study an effect of extremely early menopause in our MR study, a recent GWAS on early menopause

revealed no new genetic variants for early menopause and showed that the genetic aetiology of early menopause overlaps with that of ANM. Thus early menopause is at least partly explained by the same polygenic variants as ANM(37). Third, our analyses with glucose were based on both fasting (MAGIC) and non-fasting estimates (EPIC-CVD). Although both are associated with an increased CVD risk(38,39) it might not be appropriate to combine them, since different SNPs might drive the association and underlying mechanisms could be different.

Our findings regarding CHD are partly in contrast with one previous study investigating the association between ANM SNPs and CHD death, which reported a significantly increased risk of CHD death with a weighted genetic risk score (wGRS) in women, but not in men(17). However, our findings are in line with those of the MR analysis in women, presented in the same paper, using CARDIoGRAMplusC4D data only, which was also null. The discrepancy between the wGRS and MR findings is potentially due to the fact that the wGRS analysis was adjusted for several known CVD risk factors (current smoking, body mass index, hypertension, type 2 diabetes, total cholesterol, and lipid treatment). This might induce a biased association between the genetic variant and the outcome through confounder(s), also known as collider bias(40,41). In addition, the number of cases used for the wGRS analyses was small (only 541 CHD deaths in women), so a chance finding cannot be ruled out either.

Our MR-study suggests that the association between genetically determined reproductive aging and CHD is not causal. However, most observational studies do find an association between early age at menopause and CHD in women. We suggest several explanations for this finding. First, observational studies are susceptible to residual confounding and reverse causation. It is possible that residual confounding is still present. Postmenopausal women are by definition older than premenopausal women, making it challenging to separate the effects of

biological aging from the various phases of the reproductive aging process. Hence, residual confounding due to age may still be present in observational studies. Second, reverse causation is another potential problem in observational studies. Although most studies assume that an early ANM increases CHD risk, it might be possible that an unfavourable cardiovascular risk profile, or accelerated vascular aging, causes an early ANM. One previous study showed indeed that higher cholesterol levels prior to menopause were associated with earlier menopause(13). However, another study found no association between premenopausal CVD and subsequent age at menopause(42). If anything, women who developed CVD before menopause had a lower risk of becoming postmenopausal than women without premenopausal CVD (HR=0.98 for CVD and HR=0.90 for MI), indicating that menopause occurred later in these women(42), but none of these results were statistically significant due to the small number of premenopausal cases.

MR uses SNPs, that are randomly assigned by birth, as instrumental variables, and as such provides a method to assess causality(43). However, an MR study makes several assumptions, that have to be taken into account(44). The first assumption is that the genetic marker is associated with the exposure. The SNPs used in our study were all associated with ANM at a p-value  $<5e-6$  in the latest and largest GWAS(16). As discussed above, this may not be true in men. The second and third assumptions are that the association between the genetic marker and the outcome is explained exclusively through the exposure of interest and is unconfounded. This is often referred to as the absence of pleiotropy, which means that the genetic variant is not associated with other phenotypes. Although our Phenoscanner search showed that a few of the SNPs are associated with age at menarche or sex hormone levels, and thus that some pleiotropy may be present, our MR-Egger analysis showed no indication of

369 pleiotropy, since all intercepts were zero or very close to zero and non-significant(32). We  
370 therefore assume that our results are not biased by pleiotropy.

371         In summary, we found no evidence that reproductive aging is causally associated with  
372 CHD and CHD risk factors in women, nor in men. The association between early menopause and  
373 CHD risk in observational studies might be the result of residual confounding, reversed  
374 causation, or reflect a shared aetiology that results in both earlier menopause and higher CHD  
375 risk.

376

377

## Acknowledgements

This research has been conducted using the UK Biobank Resource under Application Number 29916. Data from m-CARDIoGRAMplusC4D have been contributed by CARDIoGRAMplusC4D investigators of the respective studies included. Statistics Netherlands is acknowledged for providing causes of death for the Dutch contribution to EPIC-CVD. We thank all EPIC participants and staff for their contribution to the study. We also thank staff from the EPIC-CVD and EPIC-InterAct Coordinating Centres for sample preparation and data handling, particularly Sarah Spackman (EPIC-CVD Data Manager) and Nicola Kerrison (EPIC-InterAct Data Manager, MRC Epidemiology Unit).

## Funding

The EPIC-CVD project was supported by the European Union Framework 7 [HEALTH-F2-2012-279233], the European Research Council [268834], the UK Medical Research Council [G0800270, MR/L003120/1], the British Heart Foundation [SP/09/002, RG/08/014, RG13/13/30194], and the UK National Institute of Health Research [to EPIC-CVD]. The establishment of the subcohort was supported by the EU Sixth Framework Programme (FP6) (grant LSHM\_CT\_2006\_037197 to the InterAct project) and the Medical Research Council Epidemiology Unit (grants MC\_UU\_12015/1 and MC\_UU\_12015/5). The national EPIC cohorts are supported by Danish Cancer Society (Denmark); Ligue Contre le Cancer, Institut Gustave Roussy, Mutuelle Générale de l'Éducation Nationale, Institut National de la Santé et de la Recherche Médicale (INSERM) (France); Deutsche Krebshilfe, Deutsches Krebsforschungszentrum and Federal Ministry of Education and Research (Germany); Ministry of Health and Social Solidarity, Stavros Niarchos Foundation and Hellenic Health Foundation

(Greece); Italian Association for Research on Cancer (AIRC) and National Research Council (Italy) and MIUR "Dipartimenti di Eccellenza"(Project D15D18000410001) to the Department of Medical Sciences; Dutch Ministry of Public Health, Welfare and Sports (VWS), LK Research Funds, Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research Fund (WCRF); ERC-2009-AdG 232997 and Nordforsk, Nordic Centre of Excellence programme on Food, Nutrition and Health (Norway); Health Research Fund (FIS), Regional Governments of Andalucía, Asturias, Basque Country, Murcia and Navarra, ISCIII RETIC [RD06/0020] (Spain); Swedish Cancer Society, Swedish Scientific Council and Regional Government of Skåne and Västerbotten (Sweden); Cancer Research UK, Medical Research Council (United Kingdom). LIFE-Heart is funded by the Leipzig Research Center for Civilization Diseases (LIFE). LIFE is an organizational unit affiliated to the Medical Faculty of the University of Leipzig. LIFE is funded by means of the European Union, by the European Regional Development Fund (ERDF) and by funds of the Free State of Saxony within the framework of the excellence initiative. This work is supported by the Dutch Heart Foundation [2013T083 to VD], by a UK Medical Research Council Skills Development Fellowship [MR/P014550/1 to SAEP] and by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (204623/Z/16/Z to SB). None of the funding sources had a role in the collection, analysis, and interpretation of the data, nor in the decision to submit the article for publication.

#### **Conflict of interest statement**

Dr. Clare Oliver-Williams received £1000 in prize money from Novartis. None of the other authors report any potential conflict of interest.



## References

1. Timmis A, Townsend N, Gale C, Grobbee R, Maniadakis N, Flather M, et al. European Society of Cardiology: Cardiovascular disease statistics 2017. *Eur Heart J*. 2018;39(7):508–77.
2. van der Schouw YT, van der Graaf Y, Steyerberg EW, Eijkemans JC, Banga JD. Age at menopause as a risk factor for cardiovascular mortality. *Lancet*. 1996;347(9003):714–8.
3. Ossewaarde ME, Bots ML, Verbeek AL, Peeters PH, van der Graaf Y, Grobbee DE, et al. Age at menopause, cause-specific mortality and total life expectancy. *Epidemiology* [Internet]. 2005/06/14. 2005;16(4):556–62. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15951675>
4. Muka T, Oliver-Williams C, Kunutsor S, Laven JSE, Fauser BCJM, Chowdhury R, et al. Association of age at onset of menopause and time since onset of menopause with cardiovascular outcomes, intermediate vascular traits, and all-cause mortality: A systematic review and meta-analysis. *JAMA Cardiol*. 2016;1(7):767–76.
5. Dam V, van der Schouw YT, Onland-Moret NC, Groenwold RHH, Peters SAE, Burgess S, et al. Association of menopausal characteristics and risk of coronary heart disease: a pan-European case-cohort analysis (submitted). 2018.
6. Matthews KA, Crawford SL, Chae CU, Everson-Rose SA, Sowers MF, Sternfeld B, et al. Are changes in cardiovascular disease risk factors in midlife women due to chronological aging or to the menopausal transition? *JAmCollCardiol*. 2009;54(25):2366–73.
7. de Kat AC, Dam V, Onland-Moret NC, Eijkemans MJC, Broekmans FJM, van der

- Schouw YT. Unraveling the associations of age and menopause with cardiovascular risk factors in a large population-based study. *BMC Med* [Internet]. BMC Medicine; 2017;15(1):2. Available from: <http://bmcmmedicine.biomedcentral.com/articles/10.1186/s12916-016-0762-8>
8. Kühnert B, Nieschlag E. Reproductive functions of the ageing male. *Hum Reprod Update*. 2004;10(4):327–39.
9. Johnson SL, Dunleavy J, Gemmell NJ, Nakagawa S. Consistent age-dependent declines in human semen quality: A systematic review and meta-analysis. *Ageing Res Rev* [Internet]. Elsevier B.V.; 2015;19:22–33. Available from: <http://dx.doi.org/10.1016/j.arr.2014.10.007>
10. Ramasamy R, Chiba K, Butler P, Lamb DJ. Male biological clock: A critical analysis of advanced paternal age. *Fertil Steril* [Internet]. Elsevier Inc.; 2015;103(6):1402–6. Available from: <http://dx.doi.org/10.1016/j.fertnstert.2015.03.011>
11. Lewis BH, Legato M, Fisch H. Medical implications of the male biological clock. *J Am Med Assoc*. 2006;296(19):2369–71.
12. Vainionpää KJ, Topo P. The making of an ageing disease: The representation of the male menopause in Finnish medical literature. *Ageing Soc*. 2005;25(6):841–61.
13. Kok HS, van Asselt KM, van der Schouw YT, van der Tweel I, Peeters PHM, Wilson PWF, et al. Heart Disease Risk Determines Menopausal Age Rather Than the Reverse. *J Am Coll Cardiol*. 2006;47(10):1976–83.
14. Smith GD, Ebrahim S. “Mendelian randomization”: Can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol*. 2003;32(1):1–

- 465 22.
- 466 15. de Haan HGG, Siegerink B, van Hylckama Vlieg A. [Mendelian randomisation]. Ned  
467 Tijdschr Geneesk [Internet]. 2014;158(44):A7547. Available from:  
468 <http://www.ncbi.nlm.nih.gov/pubmed/25322353>
- 469 16. Day FR. Europe PMC Funders Group Large-scale genomic analyses link reproductive  
470 ageing to hypothalamic signaling , breast cancer susceptibility and BRCA1-mediated  
471 DNA repair. Nat Genet. 2016;47(11):1294–303.
- 472 17. Sarnowski C, Kavousi M, Isaacs S, Demerath EW, Broer L, Muka T, et al. Genetic  
473 variants associated with earlier age at menopause increase the risk of cardiovascular  
474 events in women. Menopause [Internet]. 2017;25(4):1. Available from:  
475 <http://insights.ovid.com/crossref?an=00042192-9000000000-97667>
- 476 18. Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK Biobank: An  
477 Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases  
478 of Middle and Old Age. PLoS Med. 2015;12(3):1–10.
- 479 19. Nikpay M, Goel A, Won H, Hall LM, Willenborg C, Kanoni S, et al. A comprehensive  
480 1000 Genomes-based genome-wide association meta-analysis of coronary artery disease.  
481 Nat Genet. 2016;47(10):1121–30.
- 482 20. Danesh J, Saracci R, Berglund G, Feskens E, Overvad K, Panico S, et al. EPIC-Heart: the  
483 cardiovascular component of a prospective study of nutritional, lifestyle and biological  
484 factors in 520,000 middle-aged participants from 10 European countries. Eur J Epidemiol  
485 [Internet]. 2007/02/14. 2007;22(2):129–41. Available from:

- 486 <http://www.ncbi.nlm.nih.gov/pubmed/17295097>
- 487 21. Global Lipids Genetics Consortium, Willer CJ, Schmidt EM, Sengupta S, Peloso GM,  
488 Gustafsson S, Kanoni S, et al. Discovery and Refinement of Loci Associated with Lipid  
489 Levels. *Nat Genet.* 2013;45(11):1–24.
- 490 22. Dupuis J, Bouatia-naji N, Langenberg C. Common variants at ten genomic loci influence  
491 hemoglobin A. 2010;59(December):1–27.
- 492 23. Dupuis J, Langenberg C, Prokopenko I, Saxena R, Soranzo N, Jackson AU, et al. New  
493 genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes  
494 risk. *Nat Genet [Internet].* 2010;42(Febrero):105–16. Available from:  
495 <http://www.ncbi.nlm.nih.gov/pubmed/18554561> <http://www.nature.com/doifinder/10.1038/ng.520>  
496 0.1038/ng.520
- 497 24. Collins R. What makes UK Biobank special? *Lancet [Internet].* Elsevier Ltd;  
498 2012;379(9822):1173–4. Available from: [http://dx.doi.org/10.1016/S0140-](http://dx.doi.org/10.1016/S0140-6736(12)60404-8)  
499 6736(12)60404-8
- 500 25. Welter D, MacArthur J, Morales J, Burdett T, Hall P, Junkins H, et al. The NHGRI  
501 GWAS Catalog, a curated resource of SNP-trait associations. *Nucleic Acids Res.*  
502 2014;42(D1):1001–6.
- 503 26. Staley JR, Blackshaw J, Kamat MA, Ellis S, Surendran P, Sun BB, et al. PhenoScanner: A  
504 database of human genotype-phenotype associations. *Bioinformatics.* 2016;32(20):3207–  
505 9.
- 506 27. Burgess S, Davies NM, Thompson SG. Bias due to participant overlap in two-sample

507 Mendelian randomization. *Genet Epidemiol.* 2016;(September):597–608.

508 28. Buuren S, Groothuis-Oudshoorn K. mice: Multivariate imputation by chained equations in  
509 R. *J Stat Softw.* 2011;45(3).

510 29. Rubin DB. *Multiple Imputation for Non Response in Surveys.* Wiley; 1987.

511 30. Yavorska O, Burgess S. *MendelianRandomization: an R package for performing*  
512 *Mendelian randomization analyses using summarized data.* 2017.

513 31. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian  
514 Randomization with Some Invalid Instruments Using a Weighted Median Estimator.  
515 *Genet Epidemiol.* 2016;40(4):304–14.

516 32. Burgess S, Thompson SG. Erratum to: Interpreting findings from Mendelian  
517 randomization using the MR-Egger method (*Eur J Epidemiol*, 10.1007/s10654-017-0255-  
518 x). *Eur J Epidemiol.* Springer Netherlands; 2017;32(5):391–2.

519 33. Dale CE, Fatemifar G, Palmer TM, White J, Prieto-Merino D, Zabaneh D, et al. Causal  
520 Associations of Adiposity and Body Fat Distribution with Coronary Heart Disease, Stroke  
521 Subtypes, and Type 2 Diabetes Mellitus: A Mendelian Randomization Analysis.  
522 *Circulation.* 2017;135(24):2373–88.

523 34. Team RC. R: A language and environment for statistical computing. [Internet]. Vienna,  
524 Austria: R Foundation for Statistical computing; 2016. Available from: [https://www.r-](https://www.r-project.org/)  
525 [project.org/](https://www.r-project.org/)

526 35. Tao XY, Zuo AZ, Wang JQ, Tao FB. Effect of primary ovarian insufficiency and early  
527 natural menopause on mortality: A meta-analysis. *Climacteric.* 2016;19(1):27–36.

- 528 36. Roeters Van Lennep JE, Heida KY, Bots ML, Hoek A. Cardiovascular disease risk in  
529 women with premature ovarian insufficiency: A systematic review and meta-analysis. *Eur*  
530 *J Prev Cardiol*. 2016;23(2):178–86.
- 531 37. Perry JRB, Corre T, Esko T, Chasman DI, Fischer K, Franceschini N, et al. A genome-  
532 wide association study of early menopause and the combined impact of identified variants.  
533 *Hum Mol Genet*. 2013;22(7):1465–72.
- 534 38. Levitzky YS, Pencina MJ, D’Agostino RB, Meigs JB, Murabito JM, Vasan RS, et al.  
535 Impact of Impaired Fasting Glucose on Cardiovascular Disease. The Framingham Heart  
536 Study. *J Am Coll Cardiol*. 2008;51(3):264–70.
- 537 39. Benn M, Tybjaerg-Hansen A, McCarthy MI, Jensen GB, Grande P, Nordestgaard BG.  
538 Nonfasting glucose, ischemic heart disease, and myocardial infarction: A mendelian  
539 randomization study. *J Am Coll Cardiol* [Internet]. Elsevier Inc.; 2012;59(25):2356–65.  
540 Available from: <http://dx.doi.org/10.1016/j.jacc.2012.02.043>
- 541 40. Haycock PC, Burgess S, Wade KH, Bowden J, Relton C, Smith GD. Statistical  
542 Commentary Best ( but oft-forgotten ) practices : the design , analysis , and interpretation  
543 of Mendelian randomization studies 1. *Am J Clin Nutr*. 2016;103(August):965–78.
- 544 41. Day FR, Loh PR, Scott RA, Ong KK, Perry JRB. A Robust Example of Collider Bias in a  
545 Genetic Association Study. *Am J Hum Genet* [Internet]. The American Society of Human  
546 Genetics; 2016;98(2):392–3. Available from: <http://dx.doi.org/10.1016/j.ajhg.2015.12.019>
- 547 42. Li J, Eriksson M, Czene K, Hall P, Rodriguez-Wallberg KA. Common diseases as  
548 determinants of menopausal age. *Hum Reprod*. 2016;31(12):2856–64.

- 549 43. Burgess S, Thompson SG. Mendelian Randomization: Methods for Using Genetic  
550 Variants in Causal Estimation [Internet]. ProtoView. CRC Press; 2015. 224 p. Available  
551 from:  
552 <http://search.proquest.com/docview/1685406996?accountid=9851%5Cnhttp://tf5lu9ym5n.>  
553 [search.serialssolutions.com/?ctx\\_ver=Z39.88-2004&ctx\\_enc=info:ofi/enc:UTF-](http://search.serialssolutions.com/?ctx_ver=Z39.88-2004&ctx_enc=info:ofi/enc:UTF-)  
554 [8&rft\\_id=info:sid/ProQ%3Aagriculturejournals&rft\\_val\\_fmt=info:ofi/fmt:kev:mtx:journals](http://search.serialssolutions.com/?ctx_ver=Z39.88-2004&ctx_enc=info:ofi/enc:UTF-8&rft_id=info:sid/ProQ%3Aagriculturejournals&rft_val_fmt=info:ofi/fmt:kev:mtx:journals&rft.genre=unk)  
555 [l&rft.genre=unk](http://search.serialssolutions.com/?ctx_ver=Z39.88-2004&ctx_enc=info:ofi/enc:UTF-8&rft_id=info:sid/ProQ%3Aagriculturejournals&rft_val_fmt=info:ofi/fmt:kev:mtx:journals&rft.genre=unk)
- 556 44. Vanderweele TJ, Tchetgen EJT, Kraft P. Methodological challenges in Mendelian  
557 randomization. *Epidemiology*. 2015;25(3):427–35.
- 558